

# LARGE SCALE PERIODICITY IN REDSHIFT DISTRIBUTION

Katarzyna Bajan<sup>▲</sup>, Monika Biernacka<sup>◆</sup>, Piotr Flin<sup>1</sup><sup>■</sup>, Włodzimierz Godłowski<sup>★</sup>,  
 Victor Pervushin<sup>♣</sup>, Andrey Zorin<sup>♠</sup>

<sup>▲</sup>*H. Niewodniczanski Institute of Nuclear Physics, 31-342 Krakow, ul. Radzikowskiego 152, Poland*

<sup>◆</sup>*Pedagogical University, Institute of Physics, 25-406 Kielce, ul. Swietokrzyska 15, Poland*

<sup>■</sup>*Pedagogical University, Institute of Physics, 25-406 Kielce, ul. Swietokrzyska 15, Poland;  
 Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, 141980, Dubna, Russia*

<sup>★</sup>*Astronomical Observatory of the Jagiellonian University, 30-244 Krakow, ul. Orla 171, Poland*

<sup>♣</sup>*Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, 141980, Dubna, Russia*

<sup>♠</sup>*Faculty of Physics, MSU, Vorobjovy Gory, Moscow, 119899, Russia*

*Received October 23, 2003*

We review the previous studies of galaxies and quasar redshifts discretisation. We present also the investigations of the large scale periodicity, detected by pencil-beam observations, which revealed 128 (1/h) Mpc period, afterwards confirmed with supercluster studies. We present the theoretical possibility of obtaining such a periodicity using a toy-model. We solved the Kepler problem, i.e. the equation of motion of a particle with null energy moving in the uniform, expanding Universe, described by FLRW metrics. It is possible to obtain theoretically the separation between large scale structures similar to the observed one.

## 1. Introduction

In the large scale Universe the search of regularities is connected with testing, if radial velocities of galaxies can admit an arbitrary values, or some regular patterns, called periodisation or quantisation of galaxy redshifts is observed. The search for possible periodicity in redshift distribution has been an important question from the point of view of selection effects in observational data, as well as the correctness of applied statistics. The possible interpretation of such effect is very important too. The lack of a known mechanism causing redshift discretisation constituted the basis for claims invoking new physics at work. Due to different methods of investigations we describe separately investigations of galaxy and quasar redshift periodisation. There are presented in sections two and three. The quasar redshift periodisation in more details is presented elsewhere (Bajan et al. 2004) Moreover, the fourth section is devoted to the large scale periodicity with a period of a 128 (1/h) Mpc. This large scale periodicity obtained originally by pencil beam observations has been confirmed by various studies. So, in reality, it is the only redshift discretisation, which is almost commonly accepted. In the last section we give the possible solution of the large scale periodicity. We obtain the exact solution of the Kepler problem in the expanding Universe.

With appropriate fine tuning it is possible to obtain the distribution of structures with about 130 Mpc separation between them.

## 2. The discretisation of galaxy redshifts

The discussion of redshifts quantisation started with works of Tifft. In the Coma cluster of galaxies (Tifft 1976) optically determined redshifts of member objects can be grouped into bands with the differences of  $72.46 \text{ km} \cdot \text{s}^{-1}$ . The repetition of the analysis based on the more accurate, this is HI, data confirmed this periodicity.

The result was preceded by the study of the distribution of galaxy redshifts versus nuclear magnitudes in the Coma cluster core (Tifft 1974 and references therein), where strong redshifts periodicity of  $220 \text{ km} \cdot \text{s}^{-1}$  were observed. The existence of band structure on nuclear magnitude-redshift diagram was reported by Nanni et al. 1981 on the basis of better nuclear magnitude and redshift determination. However, in order to find the effect of quantisation it was necessary to apply to galaxy heliocentric redshift also the correction connected with the solar galactocentric movement, which is not precisely known.

Few years later Tifft and Cocke (1984) generalised

<sup>3</sup>e-mail: sfflin@cyf-kr.edu.pl

their claim for all galaxies. There exists global periodicity. It is equal to  $36.3 \text{ km} \cdot \text{s}^{-1}$  for broad-line galaxies and  $24.2 \text{ km} \cdot \text{s}^{-1}$  for narrow-line galaxies. The later value afterwards was not confirmed, so presently the value of about 36 (or 72)  $\text{km} \cdot \text{s}^{-1}$  is considered as the basic one.

Tift (1980) shown that the redshift differences for double galaxies have a statistically significant grouping near multiples of  $72 \text{ km} \cdot \text{s}^{-1}$ . Afterwards, the initial sample of galaxies has been enlarged and objects with more accurate redshift determination ( $\sigma < 25 \text{ km} \cdot \text{s}^{-1}$ ) considered (Tift 1982a,b). The periodicity of  $72 \text{ km} \cdot \text{s}^{-1}$  was observed only for such well determined redshifts. Greater uncertainties blurred the observed periodicity. The way, in which statistical analysis has been performed was criticised by Newman, Haynes & Terzian (1989). Cocke and Tift (1991) re-addressed this problem, concluding that the approach was statistically correct. Schneider and Salpeter (1991) used the sample of 107 isolated binary galaxies with very reliable 21 cm redshifts. They do not find the multiplicity of 144, 216 ...  $\text{km} \cdot \text{s}^{-1}$ , suggested by Tift, but they confirm the excess in velocity differences near  $72 \text{ km} \cdot \text{s}^{-1}$ .

Considering 40 members of the Local Group Rudnicki et al (2001) claimed the existence, at 95 % significance level, of redshifts periodisation in the considered sample, however without the precise value of the periodisation. They concluded, that the uncertainties in the published redshift determination do not allowed to make more precise statements. It is worthwhile to note, that the discretisation of redshift in the Local Group concern not only galaxies, but also globular clusters. The further analysis of the problem (Godłowski et al. 2003) shown that the strict quantisation is excluded, but a weak evidence of period  $P$  about 36 (or 24)  $\text{km} \cdot \text{s}^{-1}$  is observed.

Guthrie and Napier (1991) from the database extracted 89 non-Virgo spirals with galactocentric redshifts  $\pm 1000 \text{ km} \cdot \text{s}^{-1}$  and quoted redshift accuracy  $\sigma \leq 4 \text{ km} \cdot \text{s}^{-1}$ . An evidence of periodicity about  $37.5 \text{ km} \cdot \text{s}^{-1}$  was noted. Afterwards they (1996) repeated the analysis using 247 spirals with galactocentric radial velocities  $\pm 2600 \text{ km} \cdot \text{s}^{-1}$ , finding strong periodisation with  $P \sim 37.6 \text{ km} \cdot \text{s}^{-1}$ . The paper precisely describes the procedure of Power Spectrum Analysis, the hypothesis tested, as well as the determination of the solar vector in respect to the Galaxy centre.

### 3. The quasar redshift quantisation

This phenomenon is probably the eldest one, because in the late sixties of the previous century first claims were published. Cowan (1968) reported an effect with a period of 0.1666 in redshift  $z = \Delta\lambda/\lambda$ . The result was broadly discussed and the re-analysis of existing data, with increasing number of quasars considered,

was performed. The Power Spectrum Analysis was implemented for periodicity search, which allows to determine precisely the statistical significance of the effects (Lake & Roeder 1972). Some further claims were that periodicity is 0.089 in  $\log(1+z)$ , which corresponds to a factor 1.227 between values of two consecutive maxima in the  $1+z$  redshift distribution. Kjaergaard (1978) carefully discussed the manner, in which quasars are found and concluded, that the strong selection effects influence the observed redshift distribution, causing the existence of several peaks. So, according to his conclusions observed peaks are due to selection effects. Quite recently the sample of 1647 quasars selected in the uniform way has been analysed. Some maxima can be observed, but according to Power Spectrum Analysis they are statistically insignificant (Hawkins et al. 2002).

### 4. Large scale periodisation

Broadhurst et al. (1990) found the high peaks separated by regular spacing, strongly indicating the existence of a  $128 \cdot h^{-1} \text{ Mpc}$  (where  $h^{-1}$  is the Hubble constant in units of  $100 \text{ km} \cdot \text{s}^{-1}$ ) period in the redshift distribution. This result based on narrow pencil-beam method was confirmed by other researchers, who using the distribution of Abell galaxy clusters founded a 120 Mpc periodicity (Einasto et al. 1997). Several deep galaxy redshift surveys show this periodicity the presence of 100-Mpc scale structures (Kirshner 1997). The possibility of such large scale distribution of matter was pointed out also by Gonzales et. al (2000).

The Voronoi foam being a model describing of galaxy distribution in sheets, filaments and clusters surrounding voids was applied for study the possibility of explanation this periodicity. Van de Weygaert (1991) was able to explain the effect, but he concluded that the periodicity, even in the cellular Universe, is not a common phenomena. The similar conclusion was reached by Ikeuchi and Turner (1991), who stressed that the quasi-periodic pattern is less than only  $2\sigma$  fluctuation in a Voronoi tessellation.

The other simply possibility is the existence of a coherent peculiar velocity field with the amplitude of the order of  $10^{-3}$  and the period  $128 h^{-1} \text{ Mpc}$  (Hill, Steinhardt and Turner 1991). Such an amplitude is similar to the peculiar velocity field observed in our vicinity and connected with so-called Great Attractor. The oscillating Universe (Morikawa 1990) or the change of the values of the fundamental constants (Hill, Steinhardt and Turner (1990)) were also involved as possible explanation of this periodicity. Salgado et al. (1996), Quevedo et al. (1997) discussed the oscillation of gravitational constant.

## 5. The possible explanation of the large scale periodicity

We study the Newtonian motion of a particle in a gravitational field in the space-time with the Friedmann—Lemaitre—Robertson—Walker (FLRW) metrics

$$(ds^2) = (dt)^2 - \sum_i a^2(t)(dx^i)^2. \quad (1)$$

The observational coordinates  $X^i$  of the expanding Universe can be written as

$$X^i = a(t)x^i, \quad dX^i = a(t)dx^i + x^i da(t), \quad (2)$$

and instead of the differential of the Euclidean space  $dx^i$ , we use the covariant differential of the FLRW space coordinates

$$a(t)dx^i = d[a(t)x^i] - x^i da(t) = dX^i - X^i \frac{da(t)}{a(t)}. \quad (3)$$

One can check that the interval (1) in terms of these variables (2) becomes

$$(ds^2) = (dt)^2 - \sum_i (dX^i - H(t)X^i dt)^2, \quad (4)$$

where  $H(t) = \dot{a}(t)/a(t)$  is the Hubble parameter. As example the classical Newton action in the space with the interval (4) and the covariant derivative  $(\dot{X}^i - H(t)X^i)$  takes the following form

$$S_A = \int_{t_I}^{t_0} dt \left[ \sum_i \left( P_i (\dot{X}^i - H(t)X^i) - \frac{P_i^2}{2m_I} \right) + \frac{\alpha}{\sqrt{X_i X^i}} \right], \quad (5)$$

where  $\alpha = M_O m_I G$  is a constant of a Newtonian interaction of a galaxy with a mass  $m_I$  in a gravitational field with central mass  $M_O$ .

In the terms of the conformal time  $d\eta = dt/a$  in effective units  $r = \sqrt{x_i x^i} = R/a$ ,  $P_r = P_{Ra}$  the action (5) takes the form

$$S_A = \int_{\eta_I}^{\eta_0} d\eta \left[ P_r \frac{dr}{d\eta} + P_\theta \frac{d\theta}{d\eta} - \frac{P_r^2 + P_\theta^2/r^2}{2m} - \frac{\alpha}{r} \right], \quad (6)$$

where  $m = a(\eta)m_I$ .

Now let us consider a probably particle moving in a plane in the cylindrical coordinates

$$X^1 = R \cos \Theta, \quad X^2 = R \sin \Theta \quad (7)$$

in the gravitational field with Schwarzschild's metrics

$$ds^2 = \left( 1 - \frac{2\alpha}{mr} \right) dt^2 - \frac{dr^2}{1 - 2\alpha/(mr)} - r^2 \sin^2(\theta) d\theta^2. \quad (8)$$

In this paper we consider the case of rigid state when densities of a energy and a pressure are equals. In this case  $a(\eta) = \sqrt{1 + 2H_I(\eta - \eta_I)}$ , where the  $H_I$  is the initial value of Hubble velocity (Behnke et. al 2002). In




Figure 1: Solution equation of motion for the action (9) at  $r(\eta_I) = 1$ ,  $r'_\eta(\eta_I) = 0$ , and  $\theta(\eta_I) = 0$  for  $H_I = 1$ ,  $m_I = 1$ ,  $R_I = 1$ ,  $P_\theta = 1$ , and  $r_g = 0.5$ . For these values of initial data and integrals of motion the total system energy (defined from (9)) is equal to zero at initial point  $\eta = \eta_I$ . This point corresponds to initial point of the structure creation.

the terms of the conformal time  $d\eta = dt/a$  in relative units  $r = R/a$ ,  $P_r = P_{Ra}$  the action takes the form:

$$S_{\text{schw}} = \int_{\eta_I}^{\eta_0} d\eta \left[ P_r \frac{dr}{d\eta} + P_\theta \frac{d\theta}{d\eta} - Q_{\text{schw}} \sqrt{P_r^2 Q_{\text{schw}}^2 + P_\theta^2/r^2 + m^2 + m} \right], \quad (9)$$

where  $Q_{\text{schw}} = \left( 1 - \frac{r_g m_I}{r m} \right)^{1/2}$ ,  $r_g = M_O G$ , and  $P_r$ ,  $P_\theta$  are conjugate impulses to corresponded coordinates. The path of the probably particle is shown on the Fig. 1.

We chose the total energy of a system defined in (9) as

$$E = Q_{\text{schw}} \sqrt{P_r^2 Q_{\text{schw}}^2 + P_\theta^2/r^2 + m^2} - m \quad (10)$$

at the initial point  $\eta = \eta_I$  as equals zero. The trajectory begin at the point  $(1, 0)$ . From Fig. 5. we see that some region are more occupied then others. In such a way we show that in the expanding FLRW metrics and assuming rigid state of matter, where density equals pressure, it is possible to obtained the formation of clusters from the uniform distribution of particles with zero gravitational distributional energy. It is possible to regard this mechanism as a main reason of cellular structures is the Universe. With the fine tuning we are able to find the distributions equals to about 130 Mpc.

## 6. Conclusions

There are several effects, which are called the redshifts quantisations. The scale of this effect is drastically dif-

ferent. From the almost local values to fantastically great part of the Universe. The clear interpretation is only in the case of 100-Mpc periodisation, which is probably due to the large scale structure, but such structure origin is not easily explained. The discretisation of the quasar redshifts seems to be an artifact of selection effects and smallness of the samples. On the other hand Zhuck et al. (2001) show that the quasars seems to be located in thin walls of web with the size of  $50 \div 100$  Mpc. This finding is in good agreement with the size of voids, as well as not in great contradiction with the large scale distribution of galaxies revealed by optical data.

In the case of  $36$  (or  $72$ )  $\text{km}\cdot\text{s}^{-1}$  periodisation, which is of the main interest of the adherents of redshifts quantisation, we express the opinion that the effect is not convincingly shown. It is not quite clear, why from several thousands of galaxies in the Local Supercluster only about 250 objects (i.e. less than 4%) is taken into account, which can caused a serious bias. It will be interesting to compare the real differences between various galaxy redshifts determination and not only to consider to quoted by authors formal errors. This point was stressed also by Guthrie and Napier (1996). If this alleged effect will be confirmed, this will open totally new areas of investigations, opening also doors for so-called new physics. Beforehand, however, the detailed analysis of observational errors and applied statistical methods should be performed. The papers presenting the results have obtained till now clearly shown the weak and the strong points, which will be fruitful in further work. Moreover, the investigations of the redshift distribution in other superclusters will be of great interest. Due to accumulation of precise measurements of redshifts in the last years the re-discussion of the redshift discretisation is possible and could help to solve this bizarre problem. We show that the new physics is not needed for explanation of the existence of the large scale periodicity of about 130 Mpc. Using very simple model in which a massive particle is circulating around the source of gravitational field in the expanding Universe with FLRW metrics it is possible to obtain regions with higher density. This means that particles will be with greater probability located in some regions than others. With appropriate fine tuning it is possible to obtain the separation between dense regions (structures) of the order of 130 Mpc, which is consistent with observations.

## References

- [1] Bajan, K., Flin, P., Godłowski, W., Pervushin, V.N., El. "Particles and Nuclei", 35, 178 (2004).
- [2] D. Behnke, D.B. Blaschke, V.N. Pervushin, and D.V. Proskurin, *Phys. Lett. B*, 530, 20 (2002); [gr-qc/0102039].
- [3] Cocke, W.J., Tifft, W.G., *ApJ*, 368, 383 (1991).
- [4] Einasto, J., Einasto, M., Gottloeber, S., Mueller, V., Saar, V., Starobinsky, A.A., Tucker, D., Andern acht, H., Frisch, P., *Nature*, 385, 139 (1997).
- [5] Godłowski, W., Bajan, K., Flin, P., 2003. in preparation.
- [6] Gonzales, J.A., Quevedo, H., Salgado, M., Sudarsky, D., 2000. astro-ph/0010437.
- [7] Guthrie, B.N.G., Napier W.M., *MNRAS*, 253, 533 (1991).
- [8] Guthrie, B.N.G., Napier, W.M., *Astron. Astrophys.*, 310, 353 (1996).
- [9] Hawkins, E., Madox, S.J., Merrifield, M.N., *MNRAS*, 336, L13 (2002).
- [10] Hill, C.T., Steinhardt, P.J., Turner, M. S., *ApJ*, 366, L60 (1991).
- [11] Ikeuchi, S., Turner, E.L., *MNRAS*, 250, 519 (1991).
- [12] Kirshner, R., *Nature*, 385, 122 (1997).
- [13] Kjaergaard, P., *Phusica Scripta*, 17, 347 (1978).
- [14] Lake, R.G., Roeder, R.C., *RASC Jour.*, 66, 111 (1972).
- [15] Morikawa, M., *ApJ*, 362, L37 (1990).
- [16] Nanni, D., Pittella, G., Trevese, D., Vignato, A., *Astron. Astrophys.*, 95, 188 (1981).
- [17] Newman, W.I., Haynes, M.P., Terzian, Y., *ApJ*, 344, 111 (1989).
- [18] Quevedo, H., Salgado, M., Sudarsky, D., *GRG*, 31, 767 (1999).
- [19] Rudnicki, K., Godłowski, W., Magdziarz, A., 2001. In: "Gravitation, Electromagnetism and Cosmology: Toward a New Synthesis," (ed. K. Rudnicki) Apeiron, Montreal.
- [20] Selgado, M., Sudarsky, Quevedo, H., *Phys. Rev. D*, 53, 6771 (1996).
- [21] Schneider, S.S., Salpater, E.E., 385, 32 (1992).
- [22] Tifft W.G. 1974 in: "The formation and Dynamics of Galaxies," (J.R. Shakeshaft ed), D. Reider, Dordrecht, p. 243.
- [23] Tifft, W.G., *ApJ*, 206, 38 (1976).
- [24] Tifft, W.G., *ApJ*, 236, 70 (1980).
- [25] Tifft, W.G., *ApJ*, 257, 442 (1982a).
- [26] Tifft, W.G., *ApJ*, 262, 44 (1982b).
- [27] Tifft, W.G., Cocke, W.J., *ApJ*, 287, 492 (1984).
- [28] van de Weygaert, R., *Astron. Astrophys.*, 249, 159 (1991).
- [29] Zhuck, N.A., Moroz, V.V., Varaksin, A.M., *Spacetime & Substance*, 2, 193 (2001).